PROPOSITIONAL LOGIC II

Proof Theory

- What is logic used for? A number of things, but most importantly, it is a *language for representing the properties of things*.
- But also, we hope it will give us a *method for establishing the properties of things*.
- To see how logic may be used to establish the properties of things, it helps to look at its history.
- Logic was originally developed to make the notion of an *argument* precise.

(We do *not* mean argument as in fighting here!)

• Here is a classic argument:

All men are mortal Socrates is a man Socrates is mortal

- This example serves to illustrate a number of features of arguments:
 - The argument has a number of *premises* these are the statements that appear *before* the horizontal line;
 - The argument has a *conclusion* this is the statement that appears *after* the horizontal line;
 - The argument has the form

If

you accept that the premises are true **then** you must accept that the conclusion is true.

- In mathematics, we are concerned with when arguments are *sound*.
- To formalise the notion of a sound argument, we need some extra terminology...
- **Definition:** If $\phi \in W$, then:
 - 1. if there is *some* interpretation π such that

 $\pi \models \phi$

then ϕ is said to be *satisfiable*, otherwise ϕ is *unsatisfiable*. 2. if

 $\pi \models \phi$

for *all* interpretations π , then ϕ is said to be *valid*.

• Valid formulae of propositional logic are called *tautologies*.

• Theorem:

- 1. If ϕ is a valid formula, then $\neg \phi$ is unsatisfiable;
- 2. If $\neg \phi$ is unsatisfiable, then ϕ is valid.
- We indicate that a formula ϕ is valid by writing

 $\models \phi$.

• We can now define the *logical consequence*.

• **Definition:** If

$$\{\phi_1,\ldots,\phi_n,\phi\}\subseteq\mathcal{W}$$

then ϕ is said to be a *logical consequence* of $\{\phi_1, \ldots, \phi_n\}$ iff ϕ is satisfied by all interpretations that satisfy

 $\phi_1 \wedge \cdots \wedge \phi_n$.

• We indicate that ϕ is a logical consequence of ϕ_1, \ldots, ϕ_n by writing

$$\{\phi_1,\ldots,\phi_n\}\models\phi.$$

• An expression like this is called a *semantic sequent*.

• Theorem:

$$\{\phi_1,\ldots,\phi_n\}\models\phi.$$

iff

$$\models (\phi_1 \wedge \cdots \wedge \phi_n) \Rightarrow \phi.$$

- So we have a method for determining whether ϕ is a logical consequence of ϕ_1, \ldots, ϕ_n : we use a truth table to see whether $\phi_1 \wedge \cdots \wedge \phi_n \Rightarrow \phi$ is a tautology. If it is, then ϕ is a logical consequence of ϕ_1, \ldots, ϕ_n .
- Our main concern in proof theory is thus to have a technique for determining whether a given formula is valid, as this will then give us a technique for determining whether some formula is a logical consequence of some others.

• EXAMPLE. Show that

$$p \wedge q \models p \lor q$$
.

To do this, we construct a truth-table for

$$(p \wedge q) \Rightarrow (p \lor q).$$

Here it is:

$$p$$
 (1) (2) p q $p \land q$ $p \lor q$ F F F T F T F T T F F T T F T T

Since

$$(p \wedge q) \Rightarrow (p \vee q).$$

is true under every interpretation, we have that $p \lor q$ is a logical consequence of $p \land q$.

- The notion of logical consequence we have defined above is acceptable for a *definition* of a sound argument, but is not very helpful for checking whether a particular argument is sound or not.
- The problem is that we must look at all the possible interpretations of the primitive propositions. While this is acceptable for, say, 4 primitive propositions, it will clearly be unacceptable for 100 propositions, as it would mean checking 2¹⁰⁰ interpretations.

(Moreover, for first-order logic, there will be an *infinite* number of such interpretations.)

• What we require instead is an alternative version of logical consequence, that does not involve this kind of checking. This leads us to the idea of *syntactic* proof.

'Syntactic' Proof

- The idea of syntactic proof is to replace the semantic checking to determine whether a formula is valid by a procedure that involves purely *syntactic* manipulation.
- The kinds of techniques that we shall use are similar to those that we use when solving problems in algebra.

The basic idea is that to show that φ is a logical consequence of φ₁,..., φ_n, we use a set of *rules* to manipulate formulae.
If we can derive φ from φ₁,..., φ_n by using these rules, then φ is said to be *proved* from φ₁,..., φ_n, which we indicate by writing

$$\phi_1,\ldots,\phi_n\vdash\phi.$$

- The symbol ⊢ is called the *syntactic turnstile*.
- An expression of the form

$$\phi_1,\ldots,\phi_n\vdash\phi.$$

is called a *syntactic sequent*.

• A rule has the general form:

 $\frac{\vdash \phi_1; \cdots; \vdash \phi_n}{\vdash \phi}$ rule name

Such a rule is read:

If ϕ_1, \ldots, ϕ_n are proved then ϕ is proved.

• EXAMPLE. Here is an example of such a rule:

$$\frac{\vdash \phi; \vdash \psi}{\vdash \phi \land \psi} \land \textbf{-I}$$

This rule is called *and introduction*. It says that if we have proved ϕ , and we have also proved ψ , then we can prove $\phi \wedge \psi$.

• EXAMPLE. Here is another rule:

$$\frac{\vdash \phi \land \psi}{\vdash \phi; \vdash \psi} \land \textbf{-E}$$

This rule is called *and elimination*. It says that if we have proved $\phi \land \psi$, then we can prove both ϕ and ψ ; it allows us to eliminate the \land symbol from between them.

Let us now try to define precisely what we mean by *proof*.
Definition: (Proof) If

$$\{\phi_1,\ldots,\phi_m,\phi\}\subseteq\mathcal{W}$$

then there is a proof of ϕ from ϕ_1, \ldots, ϕ_m iff there exists some sequence of formulae

 ψ_1,\ldots,ψ_n

such that $\psi_n = \phi$, and each formula ψ_k , for $1 \le k < n$ is either one of the formula ϕ_1, \ldots, ϕ_m , or else is the conclusion of a rule whose antecedents appeared earlier in the sequence.

• If there is a proof of ϕ from ϕ_1, \ldots, ϕ_m , then we indicate this by writing:

$$\phi_1,\ldots,\phi_m\vdash\phi.$$

- It should be clear that the symbols ⊢ and ⊨ are related. We now have to state exactly *how* they are related.
- There are two properties of ⊢ to consider:
 - soundness;
 - *– completness*.
 - Intuitively, ⊢ is said to be *sound* if it is correct, in that it does not let us derive something that is not true.
 - Intuitively, *completeness* means that ⊢ will let us prove anything that is true.

• **Definition:** (Soundness) A proof system ⊢ is said to be *sound* with respect to semantics ⊨ iff

$$\phi_1,\ldots,\phi_n\vdash\phi$$

implies

$$\phi_1,\ldots,\phi_n\models\phi.$$

• **Definition:** (Completeness) A proof system ⊢ is said to be *complete* with respect to semantics ⊨ iff

$$\phi_1,\ldots,\phi_n\models\phi$$

implies

$$\phi_1,\ldots,\phi_n\vdash\phi.$$

A Proof System

- There are many proof systems for propositional logic; we shall look at a simple one.
- First, we have an unusual rule that allows us to introduce any tautology.

 $--- \frac{\text{TAUT}}{\vdash \phi} \text{ is a tautology}$

• Because a tautology is true there is no problem bringing it into the proof.



$$\frac{\vdash \phi \land \psi}{\vdash \phi; \vdash \psi} \land -\mathbf{E}$$

$$\begin{array}{c} \vdash \phi_1 \lor \cdots \lor \phi_n; \\ \phi_1 \vdash \phi; \\ \cdots; \\ \phi_n \vdash \phi \\ \vdash \phi \end{array} \lor -E$$

cis32-fall2005-parsons-lect14

18

• An alternative \lor elimination rule is:

$$\begin{array}{c} \vdash \phi \lor \psi; \\ \vdash \phi \Rightarrow \chi; \qquad \lor \mathsf{-E} \\ \hline \vdash \psi \Rightarrow \chi \\ \hline \vdash \chi \end{array}$$

• Next, a rule called *modus ponens*, which lets us eliminate \Rightarrow .

$$\frac{\vdash \phi \Rightarrow \psi; \vdash \phi}{\vdash \psi} \Rightarrow -E$$

• Next, rules for *introducing* connectives.

$$\frac{\vdash \phi_1; \cdots; \vdash \phi_n}{\vdash \phi_1 \land \cdots \land \phi_n} \land -\mathbf{I}$$

$$\frac{\vdash \phi_1; \cdots; \phi_n}{\vdash \phi_1 \lor \cdots \lor \phi_n} \lor -\mathbf{I}$$

• We have a rule called the *deduction theorem*. This rule says that if we can prove ψ from ϕ , then we can prove that $\phi \Rightarrow \psi$.

$$\frac{\phi \vdash \psi}{\vdash \phi \Rightarrow \psi} \Rightarrow -1$$

• There are a whole range of other rules, which we shall not list here.

Proof Examples

- In this section, we give some examples of proofs in the propositional calculus.
- Example 1:

$$p \wedge q \vdash q \wedge p$$

p ∧ *q* Given
 p From 1 using ∧-E
 q 1,∧-E
 q ∧ *p* 2, 3, ∧-I

• Example 2: $p \land q \vdash p \lor q$ 1. $p \wedge q$ Given 2. *p* 1, ∧-E 3. $p \lor q$ 2, \lor -I

• Example 3: $p \land q, p \Rightarrow r \vdash r$

1.
$$p \land q$$
 Given
2. p 1, \land -E
3. $p \Rightarrow r$ Given
4. r 2, 3, \Rightarrow -E

• Example 4: $p \Rightarrow q, q \Rightarrow r \vdash p \Rightarrow r$ 1. $p \Rightarrow q$ Given 2. $q \Rightarrow r$ Given 3. p Ass | 4. q 1, 3, \Rightarrow -E | 5. r 2, 4, \Rightarrow -E | 6. $p \Rightarrow r$ 3, 5, \Rightarrow -I

• Example 5: $(p \land q) \Rightarrow r \vdash p \Rightarrow (q \Rightarrow r)$

1.
$$(p \land q) \Rightarrow r$$
Given2. p Ass3. q Ass4. $p \land q$ 2, 3, \land -I5. r 1, 4, \Rightarrow -I6. $q \Rightarrow r$ 3–5, \Rightarrow -I7. $p \Rightarrow (q \Rightarrow r)$ 2–6, \Rightarrow -I

 $cis 32\mbox{-}fall 2005\mbox{-}parson s\mbox{-}lect 14$

• Example 6: $p \Rightarrow (q \Rightarrow r) \vdash (p \land q) \Rightarrow r$

1.
$$p \Rightarrow (q \Rightarrow r)$$
 Given
2. $p \land q$ Ass
3. p 2, \land -E
4. q 2, \land -E
5. $q \Rightarrow r$ 1, 3, \Rightarrow -E
6. r 4, 5, \Rightarrow -E
7. $(p \land q) \Rightarrow r$ 2-6, \Rightarrow -I

• Example 7:

$$p \Rightarrow q, \neg q \vdash \neg p$$

1.
$$p \Rightarrow q$$
 Given
2. $\neg q$ Given
3. p Ass |
4. q 1, 3, \Rightarrow -E |
5. $q \land \neg q$ 2, 4, \land -I |
6. $\neg p$ 3, 5, \neg -I

• Example 8: $p \Rightarrow q \vdash \neg (p \land \neg q)$

1.
$$p \Rightarrow q$$
Given2. $p \land \neg q$ Ass3. p 2, \land -E4. $\neg q$ 2, \land -E5. q 1, 3, \Rightarrow -E6. $q \land \neg q$ 4, 5, \land -I7. $\neg(p \land \neg q)$ 6, \neg -I

• Example 9:

Jim will party all night and pass AI? That must be wrong. If he works hard he won't have time to party. If he doesn't work hard he's not going to pass AI.

Let:

- *p* Jim will party all night*q* Jim will pass AI*r* Jim works hard
- *r* Jim works hard

Formalisation of argument:

$$r \Rightarrow \neg p, \neg r \Rightarrow \neg q \vdash \neg (p \land q)$$

1.
$$r \Rightarrow \neg p$$
 Given
2. $\neg r \Rightarrow \neg q$ Given
3. $p \land q$ Ass ||
4. r Ass ||
5. $\neg p$ 1, 4, \Rightarrow -E ||
6. p 3, \land -I ||
7. $p \land \neg p$ 5, 6, \land -I ||
8. $\neg r$ 4, 7, \neg -I |
9. $\neg q$ 2, 9, \Rightarrow -E |
10. q 3, \land -E ||
11. $q \land \neg q$ 9, 10, \land -I ||
12. $\neg (p \land q)$ 3, 11, \neg -I

Summary

- This lecture continued our look at propositional logic.
- It concentrated on proof theory, and gave examples of a number of different kinds of proof.
- Next lecture we will go on to look at predicate logic.